Numerical Linear Algebra Assignment 1

Exercise 1. (TreBau Exercise 2.5, 10 points)

Let $\mathbf{S} \in \mathbb{C}^{m \times m}$ be skew-hermitian, i.e., $\mathbf{S}^* = -\mathbf{S}$.

- (a) Show that the nonzero eigenvalues of **S** are pure imaginary.
- (b) Show that $\mathbf{I} \mathbf{S}$ is nonsingular.
- (c) Show that the matrix $\mathbf{Q} = (\mathbf{I} \mathbf{S})^{-1}(\mathbf{I} + \mathbf{S})$, know as the *Cayley transform* of **S**, is unitary. (This is a matrix analogue of a linear fractional transformation (1 + z)/(1 z), which maps the left half of the complex z-plane conformally onto the unit disk.)

Exercise 2. (TreBau Exercise 2.6, 10 points)

If **u** and **v** are *m*-vectors, the matrix $\mathbf{A} = \mathbf{I} + \mathbf{uv}^*$ is know as a rank-one perturbation of the identity. Show that if **A** is nonsingular, then its inverse has the form $\mathbf{A}^{-1} = \mathbf{I} + \alpha \mathbf{uv}^*$ for some scalar α , and give an expression for α . For what **u** and **v** is **A** singular? If it is singular, what is null(**A**)?

Exercise 3. (10 points)

Prove the Cauchy–Schwarz inequality: For any given inner product $\langle \cdot, \cdot \rangle$,

$$|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} \sqrt{\langle \mathbf{y}, \mathbf{y} \rangle}.$$

The equality holds if and only if \mathbf{x} and \mathbf{y} are linearly dependent.

Exercise 4. (10 points)

Prove that

$$\langle \mathbf{A}, \mathbf{B} \rangle = \sum_{i=1}^{m} \sum_{j=1}^{n} a_{ij} \overline{b_{ij}}, \quad \forall \mathbf{A}, \mathbf{B} \in \mathbb{C}^{m \times n}$$

is an inner product on $\mathbb{C}^{m \times n}$. (The Frobenius norm on $\mathbb{C}^{m \times n}$ is induced by this inner product.)

Exercise 5. (10 points)

Let $\|\cdot\|$ denote any vector norm on \mathbb{C}^m and $\mathbf{W} \in \mathbb{C}^{m \times m}$ be nonsingular. Prove that $\|\mathbf{x}\|_{\mathbf{W}} = \|\mathbf{W}\mathbf{x}\|$ is a vector norm on \mathbb{C}^m .

Exercise 6. (TreBau Exercise 3.2, 10 points)

Let $\|\cdot\|$ denote any vector norm on \mathbb{C}^m and also the induced matrix norm on $\mathbb{C}^{m \times m}$. Show that $\rho(\mathbf{A}) \leq \|\mathbf{A}\|$, where $\rho(\mathbf{A})$ is the *spectral radius* of \mathbf{A} , i.e., the largest absolute value $|\lambda|$ of an eigenvalue λ of \mathbf{A} .

Exercise 7. (TreBau Exercise 3.6, 10 points)

Let $\|\cdot\|$ denote any vector norm on \mathbb{C}^m . The corresponding *dual norm* $\|\cdot\|'$ is defined by the formula $\|\cdot\|' = \sup_{\|\mathbf{y}\|=1} |\mathbf{y}^*\mathbf{x}|$.

- (a) Prove that $\|\cdot\|'$ is a norm.
- (b) Let $\mathbf{x}, \mathbf{y} \in \mathbb{C}^m$ with $\|\mathbf{x}\| = \|\mathbf{y}\| = 1$ be given. Show that there exists a rank-one matrix $\mathbf{B} = \mathbf{y}\mathbf{z}^*$ such that $\mathbf{B}\mathbf{x} = \mathbf{y}$ and $\|\mathbf{B}\| = 1$, where $\|\mathbf{B}\|$ is the matrix norm of \mathbf{B} induced by the vector norm $\|\cdot\|$. You may use the following lemma, without proof: given $\mathbf{x} \in \mathbb{C}^m$, there exists a nonzero $\mathbf{z} \in \mathbb{C}^m$ such that $|\mathbf{z}^*\mathbf{x}| = \|\mathbf{z}\|'\|\mathbf{x}\|$.

Exercise 8. (10 points)

Let $\mathbf{A} \in \mathbb{C}^{m \times n}$, $\mathbf{B} \in \mathbb{C}^{n \times r}$ and let $\|\cdot\|_{\alpha}$, $\|\cdot\|_{\beta}$, and $\|\cdot\|_{\gamma}$ be norms on \mathbb{C}^{m} , \mathbb{C}^{n} , and \mathbb{C}^{r} , respectively. Prove the induced matrix norms $\|\cdot\|_{\alpha,\gamma}$, $\|\cdot\|_{\alpha,\beta}$, and $\|\cdot\|_{\beta,\gamma}$ satisfy $\|\mathbf{AB}\|_{\alpha,\gamma} \leq \|\mathbf{A}\|_{\alpha,\beta} \|\mathbf{B}\|_{\beta,\gamma}$.

Exercise 9. (10 points)

Prove that $\|\mathbf{A}\|_{\infty,1} = \max_{i,j} |a_{ij}|.$

Exercise 10. (TreBau Exercise 3.4, 10 points)

Let **A** be an $m \times n$ matrix and let **B** be a submatrix of **A**, that is, an $s \times t$ matrix ($s \le m, t \le n$) obtained by selecting certain rows and columns of **A**.

- (a) Explain how **B** can be obtained by multiplying **A** by certain row and column "deletion matices" as in step 7 of Exercise 1.1.
- (b) Using this product, show that $\|\mathbf{B}\|_p \leq \|\mathbf{A}\|_p$ for any p with $1 \leq p \leq \infty$.